

Slide 47



The content presented in this section is based on a variety of sources. In order to apply safe food handling practices, a foodservice manager needs to understand the biological, chemical, and physical hazards that can cause foodborne illness.

foodsafety

## Biological Hazards

- Seafood Toxins**
  - Ciguatera toxin
  - Scombroid toxin
  - Shellfish toxins
  - Systemic fish toxins
- Plant Toxins**
  - Poisonous plants
- Fungal Toxins**
  - poisonous mushrooms



Other Hazards 48

### FOUR SEAFOOD TOXINS (Ciguatera, Shellfish Toxin, Scombroid Toxin, and Tetrodotoxin)

#### 1. CIGUATERA POISONING

Ciguatera is a form of human poisoning caused by eating subtropical and tropical marine finfish which have accumulated naturally occurring toxins through their diet. The toxins are known to originate from several dinoflagellate (algae) species that are common to ciguatera endemic regions in the lower latitudes.

**Nature of Disease:** Manifestations of ciguatera in humans usually involves a combination of gastrointestinal, neurological, and cardiovascular disorders. Symptoms defined within these general categories vary with the geographic origin of toxic fish.

**Associated Foods:** Marine finfish most commonly implicated in ciguatera fish poisoning include the groupers, baracudas, snappers, jacks, mackerel, and triggerfish. Many other species of warm-water fishes harbor ciguatera toxins. The occurrence of toxic fish is sporadic, and not all fish of a given species or from a given locality will be toxic.

**Frequency of Disease:** The relative frequency of ciguatera fish poisoning in the United States is not known. The disease has only recently become known to the general medical community, and there is a concern that incidence is largely under-reported because of the generally non-fatal nature and short duration of the disease.

**Course of Disease and Complications:** Initial signs of poisoning occur within six hours after consumption of toxic fish and include perioral numbness and tingling (paresthesia), which may spread to the extremities, nausea, vomiting, and diarrhea. Neurological signs include intensified paresthesia, arthralgia, myalgia, headache, temperature sensory reversal and acute sensitivity to temperature extremes, vertigo, and muscular weakness to the point of prostration. Cardiovascular signs include arrhythmia, bradycardia or tachycardia, and reduced blood pressure. Ciguatera poisoning is usually self-limiting, and signs of poisoning often subside within several days from onset. However, in severe cases the neurological symptoms are known to persist from weeks to months. In a few isolated cases neurological symptoms have persisted for several years, and in other cases recovered patients have experienced recurrence of neurological symptoms months to years after recovery. Such relapses are most often associated with changes in dietary habits or with consumption of alcohol. There is a low incidence of death resulting from respiratory and cardiovascular failure.

**Target Populations:** All humans are believed to be susceptible to ciguatera toxins. Populations in tropical/subtropical regions are most likely to be affected because of the frequency of exposure to toxic fishes. However, the increasing per capita consumption of fishery products coupled with an increase in interregional transportation of seafood products has expanded the geographic range of human poisonings.

## 2. SHELLFISH TOXINS

Ingestion of contaminated shellfish results in a wide variety of symptoms, depending upon the toxins(s) present, their concentrations in the shellfish and the amount of contaminated shellfish eaten. In the case of PSP, the effects are predominantly neurological and include tingling, burning, numbness, drowsiness, incoherent speech, and respiratory paralysis. Less well characterized are the symptoms associated with DSP, NSP, and ASP. DSP is primarily observed as a generally mild gastrointestinal disorder, i.e., nausea, vomiting, diarrhea, and abdominal pain accompanied by chills, headache, and fever. Both gastrointestinal and neurological symptoms characterize NSP, including tingling and numbness of lips, tongue, and throat, muscular aches, dizziness, reversal of the sensations of hot and cold, diarrhea, and vomiting. ASP is characterized by gastrointestinal disorders (vomiting, diarrhea, abdominal pain) and neurological problems (confusion, memory loss, disorientation, seizure, coma).

**Associated Foods:** All shellfish (filter-feeding molluscs) are potentially toxic. However, PSP is generally associated with mussels, clams, cockles, and scallops; NSP with shellfish harvested along the Florida coast and the Gulf of Mexico; DSP with mussels, oysters, and scallops, and ASP with mussels.

**Relative Frequency of Disease:** Good statistical data on the occurrence and severity of shellfish poisoning are largely unavailable, which undoubtedly reflects the inability to measure the true incidence of the disease. Cases are frequently misdiagnosed and, in general, infrequently reported. Of these toxicoses, the most serious from a public health perspective appears to be PSP. The extreme potency of the PSP toxins has, in the past, resulted in an unusually high mortality rate.

## SCOMBROID POISONING

Scombroid Poisoning (also called Histamine Poisoning) Scombroid poisoning is caused by eating foods that contain high levels of histamine and possibly other vasoactive amines and compounds. Histamine and other amines are formed by the growth of certain bacteria and the subsequent action of their decarboxylase enzymes on histidine and other amino acids in food, either during the production of a product such as Swiss cheese or by spoilage of foods, such as fishery products, particularly tuna or mahi mahi. However, any food that contains the appropriate amino acids and is subjected to certain bacterial contamination and growth may lead to scombroid poisoning when eaten.

**Nature of Disease:** Initial symptoms may include a tingling or burning sensation in the mouth, a rash on the upper body and a drop in blood pressure. Frequently, headaches and itching of the skin are encountered. The symptoms may progress to nausea, vomiting, and diarrhea and may require hospitalization, particularly in the case of elderly or impaired patients.

**Associated Foods:** Fishery products that have been implicated in scombroid poisoning include the tunas (e.g., skipjack and yellowfin), mahi mahi, bluefish, sardines, mackerel, amberjack, and abalone. Many other products also have caused the toxic effects. The primary cheese involved in intoxications has been Swiss cheese. The toxin forms in a food when certain bacteria are present and time and temperature permit their growth. Distribution of the toxin within an individual fish fillet or between cans in a case lot can be uneven, with some sections of a product causing illnesses and others not. Neither cooking, canning, or freezing reduces the toxic effect. Common sensory examination by the consumer cannot ensure the absence or presence of the toxin. Chemical testing is the only reliable test for evaluation of a product.

**Relative Frequency of Disease:** Scombroid poisoning remains one of the most common forms of fish poisoning in the United States. Even so, incidents of poisoning often go unreported because of the lack of required reporting, a lack of information by some medical personnel, and confusion with the symptoms of other illnesses. Difficulties with underreporting are a worldwide problem. In the United States from 1968 to 1980, 103 incidents of intoxication involving 827 people were reported. For the same period in Japan, where the quality of fish is a national priority, 42 incidents involving 4,122 people were recorded. Since

1978, 2 actions by FDA have reduced the frequency of intoxications caused by specific products. A defect action level for histamine in canned tuna resulted in increased industry quality control. Secondly, blocklisting of mahi mahi reduced the level of fish imported to the United States.

**Course of Disease and Complications:** The onset of intoxication symptoms is rapid, ranging from immediate to 30 minutes. The duration of the illness is usually 3 hours, but may last several days.

**Target Populations:** All humans are susceptible to scombroid poisoning; however, the symptoms can be severe for the elderly and for those taking medications such as isoniazid. Because of the worldwide network for harvesting, processing, and distributing fishery products, the impact of the problem is not limited to specific geographical areas of the United States or consumption pattern. These foods are sold for use in homes, schools, hospitals, and restaurants as fresh, frozen, or processed products.

## **TETRODOTOXIN**

Pufferfish Poisoning, Tetradon Poisoning, Fugu Poisoning

**Nature of Disease:** Fish poisoning by eating members of the order Tetraodontiformes is one of the most violent intoxications from marine species. The gonads, liver, intestines, and skin of pufferfish can contain levels of tetrodotoxin sufficient to produce rapid and violent death. The flesh of many pufferfish may not usually be dangerously toxic. Tetrodotoxin has also been isolated from widely differing animal species, including the California newt, parrotfish, frogs of the genus *Atelopus*, the blue-ringed octopus, starfish, angelfish, and xanthid crabs. The metabolic source of tetrodotoxin is uncertain. No algal source has been identified, and until recently tetrodotoxin was assumed to be a metabolic product of the host. However, recent reports of the production of tetrodotoxin/anhydrotetrodotoxin by several bacterial species, including strains of the family Vibrionaceae, *Pseudomonas sp.*, and *Photobacterium phosphoreum*, point toward a bacterial origin of this family of toxins. These are relatively common marine bacteria that are often associated with marine animals. If confirmed, these findings may have some significance in toxicoses that have been more directly related to these bacterial species.

**Associated Foods:** Poisonings from tetrodotoxin have been almost exclusively associated with the consumption of pufferfish from waters of the Indo-Pacific ocean regions. Several reported cases of poisonings, including fatalities, involved pufferfish from the Atlantic Ocean, Gulf of Mexico, and Gulf of California. There have been no confirmed cases of poisoning from the Atlantic pufferfish, *Spheroides maculatus*. However, in one study, extracts from fish of this species were highly toxic in mice. The trumpet shell *Charonia sauliae* has been implicated in food poisonings, and evidence suggests that it contains a tetrodotoxin derivative. There have been several reported poisonings from mislabelled pufferfish and at least one report of a fatal episode when an individual swallowed a California newt.

**Relative Frequency of Disease:** From 1974 through 1983 there were 646 reported cases of pufferfish poisoning in Japan, with 179 fatalities. Estimates as high as 200 cases per year with mortality approaching 50% have been reported. Only a few cases have been reported in the United States, and outbreaks in countries outside the Indo-Pacific area are rare.

**Course of Disease and Complications:** The first symptom of intoxication is a slight numbness of the lips and tongue, appearing between 20 minutes to three hours after eating poisonous pufferfish. The next symptom is increasing paraesthesia in the face and extremities, which may be followed by sensations of lightness or floating. Headache, epigastric pain, nausea, diarrhea, and/or vomiting may occur. Occasionally, some reeling or difficulty in walking may occur. The second stage of the intoxication is increasing paralysis. Many victims are unable to move; even sitting may be difficult. There is increasing respiratory distress. Speech is affected, and the victim usually exhibits dyspnea, cyanosis, and hypotension. Paralysis increases and convulsions, mental impairment, and cardiac arrhythmia may occur. The victim, although completely paralyzed, may be conscious and in some cases completely lucid until shortly before death. Death usually occurs within 4 to 6 hours, with a known range of about 20 minutes to 8 hours.

**Target Populations:** All humans are susceptible to tetrodotoxin poisoning. This toxicosis may be avoided by not eating pufferfish or other animal species containing tetrodotoxin. Most other animal species known

to contain tetrodotoxin are not usually consumed by humans. Poisoning from tetrodotoxin is of major public health concern primarily in Japan, where "fugu" is a traditional delicacy. It is prepared and sold in special restaurants where trained and licensed individuals carefully remove the viscera to reduce the danger of poisoning. Importation of pufferfish into the United States is not generally permitted, although special exceptions may be granted. There is potential for misidentification and/or mislabelling, particularly of prepared, frozen fish products.

### **PLANT TOXINS**

For more information go to: <http://aggie-horticulture.tamu.edu/plantanswers/publications/poison/poison.html> or <http://www.ansci.cornell.edu/plants/>

### **FUNGAL TOXINS**

Over 5000 species of fleshy mushrooms grow naturally in North America. The vast majority have never been tested for toxicity. It is known that about 15 species are deadly and another 60 are toxic to humans whether they are consumed raw or cooked. An additional 36 species are suspected of being poisonous, whether raw or cooked. At least 40 other species are poisonous if eaten raw, but are safe after proper cooking.

Some wild mushrooms that are extremely poisonous may be difficult to distinguish from edible species. In most parts of the country there is at least one organization that include individuals who can provide assistance with both identification and program design. Governmental agencies, universities, and mycological societies are examples of such groups. If a food establishment chooses to sell wild mushrooms, management must recognize and address the need for a sound identification program for providing safe wild mushrooms.

Additional information can be found on the California Poison Control web site:  
<http://www.calpoison.org/public/mushrooms.html>.

**PHOTOGRAPH CREDITS:** The original photo appears on the following Web site:  
[http://www.vicenzanews.it/apt\\_pro/FUNGHI/HOMEPAGE.htm](http://www.vicenzanews.it/apt_pro/FUNGHI/HOMEPAGE.htm)

foodsafety

## Chemical Hazards

**Toxic Metals**

- Lead, copper, brass, zinc, antimony, cadmium

**Cleaning Agents**

- Detergents, sanitizers, polishers, abrasive cleaners, lubricants

**Pesticides and insecticides**

**Food additives**

- Preservatives (nitrite and sulfites), flavor enhancers (MSG), nutritional additives (niacin)

Other Hazards 49

### TOXIC METALS

**Lead.** Historically, lead has been used in the formulation and/or decoration of these some types of utensils. Specifically, lead-based paints that were used to decorate utensils, such as color glazes, have caused high concentrations of lead to leach into the food they contain. Lead poisoning continues to be an important public health concern due to the seriousness of associated medical problems. Lead poisoning is particularly harmful to the young and has caused learning disabilities and medical problems among individuals who have eaten high levels. The allowable levels of lead are specific to the type of utensil, based on the average contact time and properties of the foods routinely stored in each item listed.

FDA has established maximum levels for leachable lead in ceramicware, and pieces that exceed these levels are subject to recall or other agency enforcement action. The levels are based on how frequently a piece of ceramicware is used, the type and temperature of the food it holds, and how long the food stays in contact with the piece. For example, cups, mugs and pitchers have the most stringent action level, 0.5 parts per million, because they can be expected to hold food longer, allowing more time for lead to leach. Also, a pitcher may be used to hold fruit juice. And a coffee mug is generally used every day to hold a hot acidic beverage, often several times a day.

The FDA allows use of lead glazes because they're the most durable, but regulates them tightly to ensure their safety. Commercial manufacturers employ extremely strict and effective manufacturing controls that keep the lead from leaching during use. Small potters often can't control the firing of lead glazes as well so their ceramics are more likely to leach illegal lead levels, although many do use lead-free glazes.

In 21 CFR 109.16 (The Code of Federal Regulations), FDA requires high-lead-leaching decorative ceramicware to be permanently labeled that it's not for food use and may poison food. Such items bought outside the United States may not be so labeled, potentially posing serious risk if used for food.

**Copper.** High concentrations of copper are poisonous and have caused foodborne illness. When copper and copper alloy surfaces contact acidic foods, copper may be leached into the food. Carbon dioxide may be released into a water supply because of an ineffective or nonexistent backflow prevention device between a carbonator and copper plumbing components. The acid that results from mixing water and carbon dioxide leaches copper from the plumbing components and the leachate is then transferred to beverages, causing copper poisoning. Backflow prevention devices constructed of copper and copper alloys can cause, and have resulted in, the leaching of both copper and lead into carbonated beverages.

**Brass.** Brass is an alloy of copper and zinc and contains lead which is used to combine the two elements. Historically, brass has been used for items such as pumps, pipe fitting, and goblets. All three constituents are subject to leaching when they contact acidic foods, and food poisoning has resulted from such contact.

**Galvanized Metal.** Galvanized means iron or steel coated with zinc, a heavy metal that may be leached from galvanized containers into foods that are high in water content. The risk of leaching increases with increased acidity of foods contacting the galvanized container.

### **CLEANING AGENTS, PESTICIDES AND INSECTICIDES**

Preserving food safety depends in part on the appropriate and proper storage and use of poisonous or toxic materials that are necessary to the maintenance and operation of a food establishment. Even those that are necessary can pose a hazard if they are used in a manner that contradicts the intended use of the material as described by the manufacturer on the material's label. If additional poisonous or toxic materials are present, there is an unwarranted increased potential for contamination due to improper storage (e.g., overhead spillage that could result in the contamination of food, food-contact surfaces, or food equipment) or inappropriate application.

### **FOOD ADDITIVES**

Use of unapproved additives, or the use of approved additives in amounts exceeding those allowed by food additive regulations could result in foodborne illness, including allergic reactions. For example, many adverse reactions have occurred because of the indiscriminate use of sulfites to retard "browning" of fruits and vegetables or to cause ground meat to look "redder" or fresher. The concern for misuse of additives also applies to food establishments that use sodium nitrite or other curing agents in smoking and curing operations. If this process is done incorrectly, it could cause illness or death because of excessive nitrite or because the food is insufficiently preserved.

foodsafety

## Physical Hazards

- Band-aids
- Fingernails and nail polish
- Jewelry
- Broken light bulbs
- Hair
- Metal and wood
- Chipped glass
- Broken dinnerware



Other Hazards 50

Injury can result from eating a food contaminated with a hard foreign object. Hard foreign objects are called physical hazards and can result from poor procedures applied at many points in the food chain from harvest to consumer, including those within the food establishment.

### **Main Materials of Concern as Physical Hazards and Common Sources**

Glass fixtures -- can cause cuts, bleeding; may require surgery to find or remove; sources include bottles, jars, light fixtures, utensils, gauge covers

Wood -- can cause cuts, infection, choking; may require surgery to remove; sources include fields, pallets, boxes, buildings

Stones, metal fragments – can cause choking, broken teeth, cuts, infection; may require surgery to remove; sources include fields, buildings, machinery, fields, wire, employees

Insulation -- can cause choking; long-term if asbestos; sources include building materials

Bone -- can cause choking, trauma; sources include fields, improper plant processing

Plastic -- can cause choking, cuts, infection; may require surgery to remove; sources include fields, plant packaging materials, pallets, employees

Personal effects -- can cause choking, cuts, broken teeth; may require surgery to remove; source is employees

foodsafety

## Allergens

- 6-7 million Americans have food allergies.
- Most common food allergens:
  - Milk
  - Eggs
  - Fish
  - Shellfish
  - Wheat
  - Soy
  - Peanuts and tree nuts

Other Hazards 51

A food allergy is an immune system response to a food that the body mistakenly believes is harmful. Once the immune system decides that a particular food is harmful, it creates specific antibodies to it. The next time the individual eats that food, the immune system releases massive amounts of chemicals, including histamine, in order to protect the body. These chemicals trigger a cascade of allergic symptoms that can affect the respiratory system, gastrointestinal tract, skin, or cardiovascular system. Scientists estimate that approximately 11 million Americans suffer from true food allergies. At the present time, there is no cure for food allergy. Avoidance is the only way to prevent an allergic reaction.

Although an individual could be allergic to any food, such as fruits, vegetables, and meats, they are not as common as the following eight foods which account for 90% of all food-allergic reactions: milk, egg, peanut, tree nuts (walnut, cashew, etc.), fish, shellfish, soy, wheat. For more information go to: <http://www.foodallergy.org/allergens.html>

In January 2006, all foods regulated by FDA are required to be labeled as to whether it contains one or more of the top eight food allergens.

Slide 52

The slide features a decorative header image showing a variety of fresh vegetables including white rice, green peas, red bell peppers, and yellow bell peppers. The main content area is white with the word "Activity" in a large, bold, green font, followed by "Who Am I?" in a smaller, bold, black font. At the bottom of the slide, there is a wavy orange border. In the bottom right corner of this border, the text "Other Hazards" and the number "52" are visible.

**ACTIVITY INSTRUCTIONS:** Have participants tell you what type of hazard this food, object, or situation represents.

foodsafety

## Who Am I?

**Cleaners stored near food**



**Workers with long fingernails**



Other Hazards 53

**Sanitizers and cleaners stored near food** – chemical hazard

**Workers with long fingernails** – physical hazard. If the hands are not washed properly because of the long nails, microbial hazards could also be introduced into the food.

foodsafety

## Who Am I?

**Metal from a rusty grater**



**Rings on a foodservice worker's fingers**



Other Hazards 54

**Metal from a rusty grater** – chemical hazard

**Rings on a foodservice worker's fingers** – physical hazard

foodsafety

## Who Am I?

**Poisonous mushrooms**



**Wooden utensils – the wood is pine**



Other Hazards 55

**Poisonous mushrooms** – biological hazard

**Wooden utensils** – if the wood is pine – physical hazard. Hard wood is an acceptable type of wood for use in making utensils and other foodservice objects.